

The power of typicality applied to magnetic molecules and low-dimensional quantum spin systems


Jürgen Schnack

Department of Physics – University of Bielefeld – Germany

<http://obelix.physik.uni-bielefeld.de/~schnack/>

APS March Meeting 2020 Seminar, Chemistry, Colorado State University
Fort Collins, Colorado, 3 March 2020


My new office: 500 15th Street, Denver, CO 80202



We investigate magnetic molecules

J. Schnack, Contemporary Physics **60**, 127-144 (2019)

You have got a molecule!




$$S = 60!$$

Congratulations!

Powell group: npj Quantum Materials 3, 10 (2018)


You want to build a quantum computer!



Very smart!

Wernsdorfer group: Phys. Rev. Lett. **119**, 187702 (2017)


You want to deposit your molecule!



Next generation magnetic storage!

Xue group: Phys. Rev. Lett. **101**, 197208 (2008)

You want molecular magnetocalorics!



Brilliant!


Brechin group: Angew. Chem. Int. Ed. 51, 4633 (2012)

You have got an idea about the modeling!

Heisenberg

Zeeman

$$\tilde{H} = -2 \sum_{i < j} J_{ij} \vec{s}(i) \cdot \vec{s}(j) + g \mu_B B \sum_i^N s_z(i)$$




You have to solve the Schrödinger equation!

$$\underset{\sim}{H} |\phi_n\rangle = E_n |\phi_n\rangle$$

Eigenvalues E_n and eigenvectors $|\phi_n\rangle$

- needed for spectroscopy (EPR, INS, NMR);
- needed for thermodynamic functions (magnetization, susceptibility, heat capacity);
- needed for time evolution (pulsed EPR, simulate quantum computing, thermalization).

In the end it's always a big matrix!



$$\Rightarrow \begin{pmatrix} -27.8 & 3.46 & 0.18 & \cdots \\ 3.46 & -2.35 & -1.7 & \cdots \\ 0.18 & -1.7 & 5.64 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \Rightarrow$$



$\text{Fe}_{10}^{\text{III}}$: $N = 10, s = 5/2, \dim(\mathcal{H}) = (2s + 1)^N$

Dimension=60,466,176. Maybe too big?

Can we evaluate the partition function

$$Z(T, B) = \text{tr} \left(\exp \left[-\beta \tilde{H} \right] \right)$$

without diagonalizing the Hamiltonian?

Yes, we can!



$$\begin{pmatrix} 3 & 42 & 4711 \\ 42 & 0 & 3.14 \\ 4711 & 3.14 & 8 \\ -17 & 007 & 13 \\ 1.8 & 15 & 081 \end{pmatrix}$$

1. Typicality-based estimates
2. Gd_7 and the magnetocaloric effect
3. $\text{Fe}_{10}\text{Gd}_{10}$ and a quantum critical behavior
4. FTLM for anisotropic spin models
5. Bonus: Kagome lattice antiferromagnet – Is 42 the final answer?

We are the sledgehammer team of matrix diagonalization.
Please send inquiries to jschnack@uni-bielefeld.de!

Solution I: trace estimators

$$\text{tr}(\tilde{\rho}) \approx \langle r | \tilde{Q} | r \rangle$$

$$|r\rangle = \sum_{\nu} r_{\nu} |\nu\rangle, \quad r_{\nu} = \pm 1$$

- $|\nu\rangle$ some orthonormal basis of your choice; not the eigenbasis of \tilde{Q} , since we don't know it.
- $r_{\nu} = \pm 1$ random, equally distributed. Rademacher vectors.
- Amazingly accurate, bigger (Hilbert space dimension) is better.

M. Hutchinson, Communications in Statistics - Simulation and Computation **18**, 1059 (1989).

Solution II: Krylov space representation

$$\exp[-\beta \tilde{H}] \approx \tilde{\mathbb{1}} - \beta \tilde{H} + \frac{\beta^2}{2!} \tilde{H}^2 - \dots - \frac{\beta^{N_L-1}}{(N_L-1)!} \tilde{H}^{N_L-1}$$

applied to a state $|r\rangle$ yields a superposition of

$$\tilde{\mathbb{1}}|r\rangle, \quad \tilde{H}|r\rangle, \quad \tilde{H}^2|r\rangle, \quad \dots \tilde{H}^{N_L-1}|r\rangle.$$

These (linearly independent) vectors span a small space of dimension N_L ;
it is called Krylov space.

Let's diagonalize \tilde{H} in this space!

Partition function I: simple approximation

$$Z(T, B) \approx \langle r | e^{-\beta \tilde{H}} | r \rangle \approx \sum_{n=1}^{N_L} e^{-\beta \epsilon_n^{(r)}} |\langle n(r) | r \rangle|^2$$

$$O^r(T, B) \approx \frac{\langle r | Q e^{-\beta \tilde{H}} | r \rangle}{\langle r | e^{-\beta \tilde{H}} | r \rangle}$$

- Wow!!!
- One can replace a trace involving an intractable operator by an expectation value with respect to just ONE random vector evaluated by means of a Krylov space representation???

J. Jaklic and P. Prelovsek, Phys. Rev. B **49**, 5065 (1994).




Partition function II: Finite-temperature Lanczos Method

$$Z^{\text{FTLM}}(T, B) \approx \frac{1}{R} \sum_{r=1}^R \sum_{n=1}^{N_L} e^{-\beta \epsilon_n^{(r)}} |\langle n(r) | r \rangle|^2$$

- Averaging over R random vectors is better.
- $|n(r)\rangle$ n-th Lanczos eigenvector starting from $|r\rangle$ (Rademacher vectors).
- Partition function replaced by a small sum: $R = 1 \dots 100, N_L \approx 100$.

J. Jaklic and P. Prelovsek, Phys. Rev. B **49**, 5065 (1994).

FTLM 1: ferric wheel






(1) J. Schnack, J. Richter, R. Steinigeweg, Phys. Rev. Research **2**, 013186 (2020).

(2) SU(2) & D_2 : R. Schnalle and J. Schnack, Int. Rev. Phys. Chem. **29**, 403 (2010).

(3) SU(2) & C_N : T. Heitmann, J. Schnack, Phys. Rev. B **99**, 134405 (2019)




FTLM 2: icosidodecahedron



(1) J. Schnack, J. Richter, R. Steinigeweg, Phys. Rev. Research **2**, 013186 (2020).

(2) J. Schnack and O. Wendland, Eur. Phys. J. B **78**, 535 (2010).

FTLM 3: sawtooth chain



$|J_2/J_1| = 0.45$ – near critical, $|J_2/J_1| = 0.50$ – critical.


Frustration, technically speaking, works in your favour.

(1) J. Schnack, J. Richter, R. Steinigeweg, Phys. Rev. Research **2**, 013186 (2020)

(2) J. Schnack, J. Richter, T. Heitmann, J. Richter, R. Steinigeweg, Z. Naturforsch. A (2020) accepted, arXiv:2002.00411


Gd₇ and the magnetocaloric effect

Magnetocaloric effect – Paramagnets



- Ideal paramagnet: $S(T, B) = f(B/T)$, i.e. $S = \text{const} \Rightarrow T \propto B$.
- At low T pronounced effects of dipolar interaction prevent further effective cooling.


Gd₇ – Basics



- Often magnetocaloric observables not directly measured, but inferred from Maxwell's relations.
- First real cooling experiment with a molecule.
- $\hat{H} = -2 \sum_{i < j} J_{ij} \hat{s}_i \cdot \hat{s}_j + g \mu_B B \sum_i^N s_i^z$
 $J_1 = -0.090(5)$ K, $J_2 = -0.080(5)$ K
and $g = 2.02$.
- **Very good agreement down to the lowest temperatures.**

J. W. Sharples, D. Collison, E. J. L. McInnes, J. Schnack, E. Palacios, M. Evangelisti, Nat. Commun. **5**, 5321 (2014).



Gd₇ – experiment & theory



J. W. Sharples, D. Collison, E. J. L. McInnes, J. Schnack, E. Palacios, M. Evangelisti, Nat. Commun. **5**, 5321 (2014).



◀ ▶ ⌂ □ ?

✖






Fe₁₀Gd₁₀ and quantum critical behavior





$\text{Gd}_{10}\text{Fe}_{10}$ – structure = delta chain



purple: Gd ($s = 7/2$), green: Fe ($s = 5/2$)
We will see: J_1 ferro, J_2 antiferro

A. Baniodeh *et al.*, *npj Quantum Materials* **3**, 10 (2018)





Single Molecule Magnets

Finite-temperature Lanczos Method III

$$\tilde{H} = -2 \sum_{i < j} \vec{s}_i \cdot \mathbf{J}_{ij} \cdot \vec{s}_j + \sum_i \vec{s}_i \cdot \mathbf{D}_i \cdot \vec{s}_i + \mu_B B \sum_i g_i \vec{s}_i^z$$

- Problem: for anisotropic Hamiltonians no symmetry left
→ accuracy drops (esp. for high T).
- Simple traces such as $\text{Tr}(\tilde{S}^z) = 0$ tend to be wrong for R not very big.

O. Hanebaum, J. Schnack, Eur. Phys. J. B **87**, 194 (2014)

Finite-temperature Lanczos Method IV

Employ very general symmetry (time-reversal invariance)

$$\vec{\mathcal{M}}(T, -\vec{B}) = -\vec{\mathcal{M}}(T, \vec{B})$$



Use Lanczos energy eigenvector $|n(\nu)\rangle$ and time-reversed counterpart $|\tilde{n}(\nu)\rangle$

$$|n(\nu)\rangle = \sum_{\vec{m}} c_{\vec{m}} |\vec{m}\rangle \quad , \quad |\tilde{n}(\nu)\rangle = \sum_{\vec{m}} c_{\vec{m}}^* |-\vec{m}\rangle$$

- Restores $\vec{\mathcal{M}}(T, -\vec{B}) = -\vec{\mathcal{M}}(T, \vec{B})$ and (some) traces.
- More practical: use pairs of time-reversed random vectors; still accurate.

O. Hanebaum, J. Schnack, Eur. Phys. J. B **87**, 194 (2014)

Glaser-type molecules: $\text{Mn}_6^{\text{III}}\text{Cr}^{\text{III}}$



$s = 2$, $s = 3/2$



$\dim(\mathcal{H}) = 62,500$

non-collinear easy axes

Hours compared to days, notebook compared to supercomputer!

O. Hanebaum, J. Schnack, Eur. Phys. J. B **87**, 194 (2014)

A fictitious $\text{Mn}^{\text{III}}_{12}$ – M_z vs B_z





$s = 2$
 $\dim(\mathcal{H}) = 244, 140, 625$
collinear easy axes

A few days compared to *impossible*!

O. Hanebaum, J. Schnack, Eur. Phys. J. B **87**, 194 (2014)


A fictitious $\text{Mn}^{\text{III}}_{12} - M_x$ vs B_x



No other method can deliver these curves!

O. Hanebaum, J. Schnack, Eur. Phys. J. B **87**, 194 (2014)

Effective magnetic moment of Mn₁₂-acetate



We can check DFT parameter predictions for large molecules! **Normally!**

O. Hanebaum, J. Schnack, Phys. Rev. B **92** (2015) 064424


⇒ S. Ghassemi Tabrizi, A. V. Arbuznikov, and M. Kaupp, J. Phys. Chem. A **120**, 6864 (2016).



The kagome lattice antiferromagnet

Is 42 the final answer?

Kagome lattice antiferromagnet – scientific problems





- Thermodynamic functions, in particular heat capacity and susceptibility (1)
- “Condensation” of low-lying singlets below the first triplet?
- Magnetization jump to saturation
- Thermal stability of magnetization plateaus, e.g. at $M_{\text{sat}}/3$.
- Notoriously enigmatic (2)!

(1) J. Schnack, J. Schulenburg, J. Richter, Phys. Rev. B **98**, 094423 (2018)

(2) A.M. Läuchli, J. Sudan, R. Moessner, Phys. Rev. B **100**, 155142 (2019)



Kagome 42 – magnetic properties



- Low- T peak moves to higher T with increasing N .
- Density of low-lying singlets seems to move to higher excitation energies!

J. Schnack, J. Schulenburg, J. Richter, Phys. Rev. B **98**, 094423 (2018)


Kagome – tensor network calculations



- Tensor network calculations for the infinite system (1).

(1) Xi Chen, Shi-Ju Ran, Tao Liu, Cheng Peng, Yi-Zhen Huang, Gang Su, Science Bulletin **63**, 1545 (2018).

Summary



- Magnetic molecules for storage, q-bits, MCE, and since they are nice.
- Molecules taught us about frustrated systems.
- Isentropes for interacting systems are much richer than for paramagnets. Good for applications away from $(T = 0, B = 0)$.
- Quantum phase transitions may allow barocaloric applications.
- ED, HTE, CMC, QMC, FTLM, DMRG, DDMRG, thDMRG for magnetic molecules.

Many thanks to my collaborators



- C. Beckmann, M. Czopnik, T. Glaser, O. Hanebaum, Chr. Heesing, M. Höck, N.B. Ivanov, H.-T. Langwald, A. Müller, R. Schnalle, Chr. Schröder, J. Ummethum, P. Vorndamme (Bielefeld)
- K. Bärwinkel, T. Heitmann, R. Heveling, H.-J. Schmidt, R. Steinigeweg (Osnabrück)
- M. Luban (Ames Lab); P. Kögerler (Aachen, Jülich, Ames); D. Collison, R.E.P. Winpenny, E.J.L. McInnes, F. Tuna (Man U); L. Cronin, M. Murrie (Glasgow); E. Brechin (Edinburgh); H. Nojiri (Sendai, Japan); A. Postnikov (Metz); M. Evangelisti (Zaragoza); A. Honecker (U Cergy-Pontoise); E. Garlatti, S. Carretta, G. Amoretti, P. Santini (Parma); A. Tenant (ORNL); Gopalan Rajaraman (Mumbai); M. Affronte (Modena)
- J. Richter, J. Schulenburg (Magdeburg); B. Lake (HMI Berlin); B. Büchner, V. Kataev, H.-H. Klauß (Dresden); A. Powell, W. Wernsdorfer (Karlsruhe); J. Wosnitza (Dresden-Rossendorf); J. van Slageren (Stuttgart); R. Klingeler (Heidelberg); O. Waldmann (Freiburg)

Thank you very much for your
attention.

The end.

Molecular Magnetism Web

www.molmag.de

Highlights. Tutorials. Who is who. Conferences.