

Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions

Patrick Vorndamme, Heinz-Jürgen Schmidt, Christian Schröder, Jürgen Schnack

Department of Physics – University of Bielefeld – Germany

<http://obelix.physik.uni-bielefeld.de/~schnack/>

Theory Seminar


Osnabrück University, 4 April 2022

Movie 1


(Don't relax! You will be asked what you saw!)

Synchronization I – Setting


- System of N spins (mostly $s = 1/2$);
- Unitary time evolution with Hamiltonian \tilde{H} ;
- Zeeman term included, field along z -direction;
- Initial state, e.g. product state, with single spin expectation values in x - y -plane;
- Let go!
- What do you expect?




(a) 0 degrees



(b) 180 degrees



(c) 360 degrees




(d) random directions

P. Vorndamme, H.-J. Schmidt, Chr. Schröder, J. Schnack, *Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions*, New J. Phys. **23**, 083038 (2021)

Synchronization I – Setting

- System of N spins (mostly $s = 1/2$);
- Unitary time evolution with Hamiltonian \tilde{H} ;
- Zeeman term included, field along z -direction;
- Initial state, e.g. product state, with single spin expectation values in x - y -plane;
- Let go!
- What do you expect?
Want to see the movie again?



P. Vorndamme, H.-J. Schmidt, Chr. Schröder, J. Schnack, *Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions*, New J. Phys. **23**, 083038 (2021)

Synchronization II – Heisenberg case


- $$\underline{H} = - \sum_{j=1}^N J_j \underline{\tilde{s}}_j \cdot \underline{\tilde{s}}_{j+1} - \sum_{j=1}^N h_j s_j^z \quad (1);$$

- $\forall j : h_j = h$: total spin and transverse magnetization conserved;

$$M_{\text{trans}} := \sqrt{\langle \underline{S}^x \rangle^2 + \langle \underline{S}^y \rangle^2};$$

- Not entangled: purity $\text{Tr}(\rho_j^2) = 1$,
maximally entangled: purity $\text{Tr}(\rho_j^2) = 0.5$;

- Let go with random J_j !
- What do you expect?



Time evolution of initial state $|\psi_B\rangle$ w.r.t. Hamiltonian (1) with isotropic Heisenberg interactions and $J_j \in [1.6, 2.4]$, $h_j = -1 \forall j$, $N = 25$.

Synchronization II – Heisenberg case

- $$\underline{H} = - \sum_{j=1}^N J_j \underline{\tilde{s}}_j \cdot \underline{\tilde{s}}_{j+1} - \sum_{j=1}^N h_j s_j^z \quad (1);$$


- $\forall j : h_j = h$: total spin and transverse magnetization conserved;

$$M_{\text{trans}} := \sqrt{\langle \underline{S}^x \rangle^2 + \langle \underline{S}^y \rangle^2};$$

- Not entangled: purity $\text{Tr}(\rho_j^2) = 1$,
maximally entangled: purity $\text{Tr}(\rho_j^2) = 0.5$;

- Let go with random J_j !

- What do you expect?
Want to see the movie again?



Time evolution of initial state $|\psi_B\rangle$ w.r.t. Hamiltonian (1) with isotropic Heisenberg interactions and $J_j \in [1.6, 2.4]$, $h_j = -1 \forall j$, $N = 25$.

Synchronization III – our understanding




- We understand the case where all $J_j = J$ and all $h_j = h$.
- Total spin and transverse magnetization conserved;
- AND: all spins equivalent!
- If one assumes local equilibration to a state compatible with the conserved quantities, then all spins need to have the same expectation value.
- Synchronization is observed for the vast majority of all initial states and Heisenberg Hamiltonians that we investigated so far.

What about similar systems of classical spins?

Movie 2

(Schröder Bros. Inc. – Bielefeld, Melle,
Hollywood)

Synchronization IV – classical Heisenberg case




Time evolution of initial states A, dots, D w.r.t. classical Hamiltonian (1) with isotropic Heisenberg interactions and $J_j \in [1.6, 2.4]$, $h_j = -1 \forall j$, $N = 24$.

P. Vorndamme, H.-J. Schmidt, Chr. Schröder, J. Schnack, *Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions*, New J. Phys. **23**, 083038 (2021)

- $H = - \sum_{j=1}^N J_j \vec{s}_j \cdot \vec{s}_{j+1} - \sum_{j=1}^N h_j s_j^z \quad (1);$
- Classical spins do not synchronize in a closed system. Never!
- Classical spins have N additional conserved quantities, the length of the classical spins.
- Classical spins cannot entangle.

Synchronization IV – classical Heisenberg case



Time evolution of initial states A, dots, D w.r.t. classical Hamiltonian (1) with isotropic Heisenberg interactions and $J_j \in [1.6, 2.4]$, $h_j = -1 \forall j$, $N = 24$.

P. Vorndamme, H.-J. Schmidt, Chr. Schröder, J. Schnack, *Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions*, New J. Phys. **23**, 083038 (2021)


- $H = - \sum_{j=1}^N J_j \vec{s}_j \cdot \vec{s}_{j+1} - \sum_{j=1}^N h_j s_j^z \quad (1);$
- Classical spins do not synchronize in a closed system. Never!
- Classical spins have N additional conserved quantities, the length of the classical spins.
- Classical spins cannot entangle.
Want to see the movie again?

What about other systems
in the zoo of spin Hamiltonians?

Movie 3

(Guess what happens to the purity!)


Synchronization V – loss of symmetries



Time evolution of initial state $|\psi_B\rangle$ w.r.t. Hamiltonian (2) with for two values of δ , and $N = 24$, $J = 2$, $h = -1$.

- $$\begin{aligned} \tilde{H}_{XYZ} = & -J \sum_{j=1}^N \tilde{s}_j^x \tilde{s}_{j+1}^x \\ & - (J - \delta) \sum_{j=1}^N \tilde{s}_j^y \tilde{s}_{j+1}^y \\ & - (J - 2\delta) \sum_{j=1}^N \tilde{s}_j^z \tilde{s}_{j+1}^z - h \sum_{j=1}^N \tilde{s}_j^z \quad (2); \end{aligned}$$
- Hamiltonians with less symmetries down to none;
- What do you expect?

Synchronization V – loss of symmetries



Time evolution of initial state $|\psi_B\rangle$ w.r.t. Hamiltonian (2) with for two values of δ , and $N = 24$, $J = 2$, $h = -1$.

- $$\begin{aligned} \tilde{H}_{XYZ} = & -J \sum_{j=1}^N \tilde{s}_j^x \tilde{s}_{j+1}^x \\ & - (J - \delta) \sum_{j=1}^N \tilde{s}_j^y \tilde{s}_{j+1}^y \\ & - (J - 2\delta) \sum_{j=1}^N \tilde{s}_j^z \tilde{s}_{j+1}^z - h \sum_{j=1}^N \tilde{s}_j^z \quad (2); \end{aligned}$$

- Hamiltonians with less symmetries down to none;

- What do you expect?
Transient synchronization and decay to zero!
Want to see the movie again?

P. Vorndamme, H.-J. Schmidt, Chr. Schröder, J. Schnack, *Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions*, New J. Phys. **23**, 083038 (2021)

Thank you very much for your attention.



Patrick Vorndamme



Christian Schröder



Heinz-Jürgen Schmidt



Jürgen Schnack

The end.

Magnon crystallization in the kagome lattice antiferromagnet



P. Vorndamme, H.-J. Schmidt, Chr. Schröder, J. Schnack, *Observation of phase synchronization and alignment during free induction decay of quantum spins with Heisenberg interactions*, New J. Phys. **23**, 083038 (2021)