

Exact eigenstates of highly frustrated spin lattices probed in high fields

Jürgen Schnack, H.-J. Schmidt, A. Honecker, J. Schulenburg, and J. Richter

Osnabrück – Göttingen – Magdeburg, Germany
<http://obelix.physik.uni-osnabrueck.de/~schnack/>

Yamada Conference LX on Research in High Magnetic Fields
August 16-19, 2006, Sendai, Japan



Thanks to many collaborators ...


- K. Bärwinkel, H.-J. Schmidt, J. S., M. Allalen, M. Brüger, D. Mentrup, D. Müter, M. Exler, P. Hage, F. Hesmer, K. Jahns, F. Ouchni, R. Schnalle, P. Shchelokovskyy, S. Torbrügge & M. Neumann, K. Küpper, M. Prinz (UOS);
- M. Luban, P. Kögerler, D. Vaknin (Ames Lab, USA);
J. Musfeld (U. of Tennessee, USA); N. Dalal (Florida State, USA);
- R.E.P. Winpenny (Man U, UK); L. Cronin (U. of Glasgow, UK);
H. Nojiri (Tohoku University, Japan);
- A. Müller (U. Bielefeld) & Chr. Schröder (FH Bielefeld);
J. Richter, J. Schulenburg, R. Schmidt (U. Magdeburg);
S. Blügel, A. Postnikov (FZ Jülich); A. Honecker (U. Göttingen);
E. Rentschler (U. Mainz); U. Kortz (IUB); A. Tennant, B. Lake (HMI Berlin);
- B. Büchner, V. Kataev, R. Klingeler (IFW Dresden)

... who helped to obtain various general results

1. Extension of Lieb, Schultz, and Mattis: k -rule for spin odd rings
2. Rotational bands in antiferromagnets
3. Giant magnetization jumps in frustrated antiferromagnets
4. Magnetization plateaus and susceptibility minima
5. Enhanced magnetocaloric effect
6. Hysteresis without anisotropy
7. Special properties of a triangular molecule-based spin tube

My favorite starting point

$\{\text{Mo}_{72}\text{Fe}_{30}\}$ – a molecular brother of the kagome lattice and an archetype of geometric frustration



- Giant magnetic Keplerate molecule;
- Structure: Fe - yellow, Mo - blue, O - red;
- Antiferromagnetic interaction mediated by O-Mo-O bridges (1).
- Classical ground state of $\{\text{Mo}_{72}\text{Fe}_{30}\}$: three sublattice structure, coplanar spins (2);
- Quantum mechanical ground state $S = 0$ can only be approximated, dimension of Hilbert space $(2s + 1)^N \approx 10^{23}$ (3).

(1) A. Müller *et al.*, Chem. Phys. Chem. **2**, 517 (2001) , (2) M. Axenovich and M. Luban, Phys. Rev. B **63**, 100407 (2001) , (3) M. Exler and J. Schnack, Phys. Rev. B **67**, 094440 (2003)

Hamiltonian

Model Hamiltonian – Heisenberg-Model

$$\tilde{H} = \sum_{i,j} \vec{s}(i) \cdot \mathbf{J}_{ij} \cdot \vec{s}(j) + \sum_{i,j} \vec{D}_{ij} \cdot [\vec{s}(i) \times \vec{s}(j)] + \mu_B B \sum_i^N g_i \tilde{s}_z(i)$$

Exchange/Anisotropy Dzyaloshinskii-Moriya Zeeman

Very often anisotropic terms are utterly negligible, then . . .


$$\tilde{H} = - \sum_{i,j} J_{ij} \vec{s}(i) \cdot \vec{s}(j) + g \mu_B B \sum_i^N \tilde{s}_z(i)$$

Heisenberg Zeeman

The Heisenberg Hamilton operator together with a Zeeman term are used for the following considerations; $J < 0$: antiferromagnetic coupling.

From rotational bands to giant magnetization jumps

Rotational bands in antiferromagnets I




- Often minimal energies $E_{min}(S)$ form a rotational band: Landé interval rule (1);
- For bipartite systems (2,3): $\tilde{H}^{\text{eff}} = -2 J_{\text{eff}} \tilde{\vec{S}}_A \cdot \tilde{\vec{S}}_B$;
- Lowest band – rotation of Néel vector, second band – spin wave excitations (4).

- (1) A. Caneschi *et al.*, Chem. Eur. J. **2**, 1379 (1996), G. L. Abbati *et al.*, Inorg. Chim. Acta **297**, 291 (2000)
- (2) J. Schnack and M. Luban, Phys. Rev. B **63**, 014418 (2001)
- (3) O. Waldmann, Phys. Rev. B **65**, 024424 (2002)
- (4) P.W. Anderson, Phys. Rev. B **86**, 694 (1952), O. Waldmann *et al.*, Phys. Rev. Lett. **91**, 237202 (2003).

Rotational bands in antiferromagnets II

Approximate Hamiltonian for $\{\text{Mo}_{72}\text{Fe}_{30}\}$



$$\tilde{H} = -2J \sum_{(u < v)} \tilde{s}(u) \cdot \tilde{s}(v) \approx -2J_{\text{eff}} \left[\tilde{s}_A \cdot \tilde{s}_B + \tilde{s}_B \cdot \tilde{s}_C + \tilde{s}_C \cdot \tilde{s}_A \right] = \tilde{H}^{\text{eff}}$$

Three sublattice system, classical 120° -ground state;
Good description of low-temperature magnetization.


J. Schnack, M. Luban, R. Modler, Europhys. Lett. **56**, 863 (2001)

Surprise!

The parabola is straight . . .
. . . at least at the top end!

Giant magnetization jumps in frustrated antiferromagnets I

$\{\text{Mo}_{72}\text{V}_{30}\}$



- Close look: $E_{\min}(S)$ linear in S for high S instead of being quadratic (1);
- Heisenberg model: property depends only on the structure but not on s (2);
- Alternative formulation: independent localized magnons (3);


(1) J. Schnack, H.-J. Schmidt, J. Richter, J. Schulenburg, Eur. Phys. J. B **24**, 475 (2001)

(2) H.-J. Schmidt, J. Phys. A: Math. Gen. **35**, 6545 (2002)

(3) J. Schulenburg, A. Honecker, J. Schnack, J. Richter, H.-J. Schmidt, Phys. Rev. Lett. **88**, 167207 (2002)

Giant magnetization jumps in frustrated antiferromagnets II

Localized Magnons




- $|\text{localized magnon}\rangle = \frac{1}{2}(|1\rangle - |2\rangle + |3\rangle - |4\rangle)$
 - $|1\rangle = \tilde{s}^-(1)|\uparrow\uparrow\uparrow\dots\rangle$ etc.
 - $\tilde{H}|\text{localized magnon}\rangle \propto |\text{localized magnon}\rangle$
 - Localized magnon is state of lowest energy (1,2).
-
- Triangles trap the localized magnon, amplitudes cancel at outer vertices.

- (1) J. Schnack, H.-J. Schmidt, J. Richter, J. Schulenburg, Eur. Phys. J. B **24**, 475 (2001)
(2) H.-J. Schmidt, J. Phys. A: Math. Gen. **35**, 6545 (2002)

Giant magnetization jumps in frustrated antiferromagnets III

Kagome Lattice




- Non-interacting one-magnon states can be placed on various lattices, e. g. kagome or pyrochlore;
- Each state of n independent magnons is the ground state in the Hilbert subspace with $M = Ns - n$; Kagome: max. number of indep. magnons is $N/9$;
- Linear dependence of E_{\min} on M
⇒ magnetization jump;
- Jump is a macroscopic quantum effect!
- A rare example of analytically known many-body states!

J. Schulenburg, A. Honecker, J. Schnack, J. Richter, H.-J. Schmidt, Phys. Rev. Lett. **88**, 167207 (2002)
 J. Richter, J. Schulenburg, A. Honecker, J. Schnack, H.-J. Schmidt, J. Phys.: Condens. Matter **16**, S779 (2004)

Giant magnetization jumps in frustrated antiferromagnets IV


Example: kagome chain



- Minimal energies E_{\min} of a kagome chain with $N = 36$ and $s = 1/2$.
- ($T = 0$)-magnetization curve of the kagome chain with $N = 36$. The magnetization jump is $\Delta M = 6$ (1).

(1) J. Schnack, H.-J. Schmidt, A. Honecker, J. Schulenburg, and J. Richter, cond-mat/0606401

Condensed matter physics point of view: Flat band



- Flat band of minimal energy in one-magnon space, i. e. high degeneracy of ground state energy in $\mathcal{H}(M = Ns - 1)$;
- Localized magnons can be built from those eigenstates of the translation operator, that belong to the flat band;
- There is a relation to flat band ferromagnetism (H. Tasaki & A. Mielke), compare (1).

(1) A. Honecker, J. Richter, Condens. Matter Phys. **8**, 813 (2005)
 J. Schnack, H.-J. Schmidt, A. Honecker, J. Schulenburg, and J. Richter, cond-mat/0606401

Enhanced magnetocaloric effect I

Basics

$$\left(\frac{\partial T}{\partial B}\right)_S = -\frac{T}{C} \left(\frac{\partial S}{\partial B}\right)_T$$

(adiabatic temperature change)

- Heating or cooling in a varying magnetic field. Discovered in pure iron by E. Warburg in 1881.
- Typical rates: 0.5 ... 2 K/T.
- Giant magnetocaloric effect: 3 ... 4 K/T e.g. in $\text{Gd}_5(\text{Si}_x\text{Ge}_{1-x})_4$ alloys ($x \leq 0.5$).

- MCE especially large at large isothermal entropy changes, i.e. at phase transitions (1), close to quantum critical points (2), or due to the condensation of independent magnons (3).


(1) V.K. Pecharsky, K.A. Gschneidner, Jr., A. O. Pecharsky, and A. M. Tishin, Phys. Rev. B **64**, 144406 (2001)

(2) Lijun Zhu, M. Garst, A. Rosch, and Qimiao Si, Phys. Rev. Lett. **91**, 066404 (2003)

(3) M.E. Zhitomirsky, A. Honecker, J. Stat. Mech.: Theor. Exp. **2004**, P07012 (2004)

Enhanced magnetocaloric effect II


Simple af $s = 1/2$ dimer



- Singlet-triplet level crossing leads to degenerate ground state and thus non-vanishing entropy at $T = 0$.
- $M(T = 0, B)$ and $S(T = 0, B)$ not analytic as function of B .
- $C(T, B)$ varies strongly as function of B for low T .

Enhanced magnetocaloric effect IV

Isentrops of af $s = 1/2$ dimer



blue lines: ideal paramagnet, red curves: af dimer

Magnetocaloric effect:


- (a) reduced,
- (b) the same,
- (c) enhanced,
- (d) opposite

when compared to an ideal paramagnet.

Case (d) does not occur for a paramagnet.


Enhanced magnetocaloric effect V

Two molecular spin systems



- Graphics: isentrops of the frustrated cuboctahedron and a $N = 12$ ring molecule;
- Cuboctahedron features independent magnons and extraordinarily high jump to saturation;
- Degeneracy and ($T = 0$)–entropy at saturation field higher for the cuboctahedron;
- Adiabatic (de-) magnetization more efficient for the frustrated spin system.

Discussion



- In real compounds the perfect degeneracy at B_{sat} would be lifted, e.g. by dipolar or Dzyaloshinskii-Moriya interactions.
- Magnetization jump would be smeared out.
- Low-lying density of states remains large at B_{sat} , which would still be clearly visible in magnetocaloric investigations.

Summary

Frustration can lead to exotic behavior.

And, the end is not in sight, . . .

... , however, this talk is at its end!

Thank you very much for your attention.

German Molecular Magnetism Web

www.molmag.de

Highlights. Tutorials. Who is who. DFG SPP 1137