

Exact and quasi exact numerical methods for giant magnetic molecules



Jürgen Schnack

Department of Physics – University of Bielefeld – Germany

<http://obelix.physik.uni-bielefeld.de/~schnack/>


APS March meeting, Session P13

Boston, 29. 02. 2012



The problem

You have got a molecule!




Congratulations!

You have got an idea about the modeling!


$$\tilde{H} = -2 \sum_{i < j} J_{ij} \vec{\tilde{s}}(i) \cdot \vec{\tilde{s}}(j) + g \mu_B B \sum_i^N \tilde{s}_z(i)$$

Heisenberg

Zeeman




In the end it's always a big matrix!



$\text{Fe}_{10}^{\text{III}}$: $N = 10, s = 5/2$

Dimension=60,466,176. Maybe too big?

Thank God, we have computers








“cell professor”

128 cores, 384 GB RAM

... but that's not enough!

Magnetic Molecules



possess symmetries! Use them!

Key question:


Can we employ $SU(2)$ symmetry
together with arbitrary
point group symmetries?

SU(2) symmetry

Quantum chemists need to be much smarter since they have smaller computers!

- (1) D. Gatteschi and L. Pardi, *Gazz. Chim. Ital.* **123**, 231 (1993).
- (2) J. J. Borrás-Almenar, J. M. Clemente-Juan, E. Coronado, and B. S. Tsukerblat, *Inorg. Chem.* **38**, 6081 (1999).
- (3) B. S. Tsukerblat, *Group theory in chemistry and spectroscopy: a simple guide to advanced usage*, 2nd ed. (Dover Publications, Mineola, New York, 2006).

Irreducible Tensor Operator approach



Spin rotational symmetry:

- $\underline{H} = -2 \sum_{i < j} J_{ij} \vec{s}_i \cdot \vec{s}_j + g\mu_B \vec{S} \cdot \vec{B}$;
- $[\underline{H}, \vec{S}^2] = 0, [\underline{H}, S_z] = 0$;
- Irreducible Tensor Operator (ITO) approach;
- Free program MAGPACK (2) available.

(1) D. Gatteschi and L. Pardi, Gazz. Chim. Ital. **123**, 231 (1993).

(2) J. J. Borrás-Almenar, J. M. Clemente-Juan, E. Coronado, and B. S. Tsukerblat, Inorg. Chem. **38**, 6081 (1999).

Idea of ITO

$$\begin{aligned}
 \underline{H}_{\text{Heisenberg}} &= -2 \sum_{i < j} J_{ij} \vec{s}_i \cdot \vec{s}_j \\
 &= 2\sqrt{3} \sum_{i < j} J_{ij} \underline{T}^{(0)}(\{k_i\}, \{\bar{k}_i\} | k_i = k_j = 1)
 \end{aligned}$$

Irreducible Tensor Operator approach

- Express spin operators and functions thereof as ITOs;
- Use vector coupling basis $|\alpha S M\rangle$ and recursive recoupling.

(1) Gatteschi, Tsukerblat, Coronado, Waldmann, ...
 (2) R. Schnalle, Ph.D. thesis, Osnabrück University (2009)

Point group symmetry

(Ph.D. of Roman Schnalle)

Point Group Symmetry I

$$|\alpha' S M \Gamma\rangle = \mathcal{P}^{(\Gamma)} |\alpha S M\rangle = \left(\frac{l_{\Gamma}}{h} \sum_R \left(\chi^{(\Gamma)}(R) \right)^* \mathcal{G}(R) \right) |\alpha S M\rangle$$

Method:

- *Basis function generating machine* (1);
- **Projection on irreducible representations Γ (Wigner);**
- **Orthonormalization necessary.**

(1) M. Tinkham, *Group Theory and Quantum Mechanics*, Dover.

(2) D. Gatteschi and L. Pardi, *Gazz. Chim. Ital.* **123**, 231 (1993).

(3) O. Waldmann, *Phys. Rev. B* **61**, 6138 (2000).

(4) R. Schnalle and J. Schnack, *Int. Rev. Phys. Chem.* **29**, 403-452 (2010).




Point Group Symmetry II

$$\tilde{G}(R) |\alpha S M\rangle_a = |\alpha S M\rangle_b = \sum_{\alpha'} |\alpha' S M\rangle_a {}_a\langle \alpha' S M | \alpha S M \rangle_b$$

- Serious problem: application of $\tilde{G}(R)$, i.e. permutation of spins, leads to different coupling schemes: $a \Rightarrow b$;
- Solution: implementation of graph-theoretical results to evaluate recoupling coefficients ${}_a\langle \alpha' S M | \alpha S M \rangle_b$.


Results

Results I: Cuboctahedron





Cuboctahedron, $s = 3/2$, Hilbert space dimension 16,777,216; symmetry O_h (1).
 Evaluation of recoupling coefficients very time consuming (1,2).

- (1) J. Schnack and R. Schnalle, Polyhedron **28**, 1620 (2009).
- (2) R. Schnalle and J. Schnack, Phys. Rev. B **79**, 104419 (2009).




Results II: Icosahedron





Icosahedron, $s = 3/2$, Hilbert space dimension 16,777,216; symmetry I_h ;
 Evaluation of recoupling coefficients for $s = 3/2$ in I_h **practically impossible** (1).

(1) R. Schnalle and J. Schnack, Int. Rev. Phys. Chem. **29**, 403-452 (2010).



Results III: Fe₁₀



Spin ring, $N = 10$, $s = 5/2$, Hilbert space dimension 60,466,176; symmetry D_2 ; Symmetry C_{10} would lead to more complicated recoupling coefficients & complex representation (1).

(1) R. Schnalle and J. Schnack, *Int. Rev. Phys. Chem.* **29**, 403-452 (2010).


(2) C. Delfs *et al.*, *Inorg. Chem.* **32**, 3099 (1993).

This could be the end . . .

... but you have got another molecule!

$N=30, s=1/2$

DIMENSION = 1,073,741,824



Congratulations!

The idea of finite-temperature Lanczos

$$Z(T, B) = \sum_{\nu} \langle \nu | \exp \left\{ -\beta \tilde{H} \right\} | \nu \rangle$$



$$\langle \nu | \exp \left\{ -\beta \tilde{H} \right\} | \nu \rangle \approx \sum_n \langle \nu | n(\nu) \rangle \exp \left\{ -\beta \epsilon_n \right\} \langle n(\nu) | \nu \rangle$$

$$Z(T, B) \approx \frac{\dim(\mathcal{H})}{R} \sum_{\nu=1}^R \sum_{n=1}^{N_L} \exp \left\{ -\beta \epsilon_n \right\} |\langle n(\nu) | \nu \rangle|^2$$

- $|n(\nu)\rangle$ n-th Lanczos eigenvector starting from $|\nu\rangle$;
- Partition function replaced by a small sum: $R = 1 \dots 10, N_L \approx 100$.

J. Jaklic and P. Prelovsek, Phys. Rev. B **49**, 5065 (1994).

How good is finite-temperature Lanczos?




- Works very well, see e.g. cuboctahedron and icosahedron.
- $N = 12, s = 3/2$: Considered $< 100,000$ states instead of 16,777,216.

Exact results: R. Schnalle and J. Schnack, Int. Rev. Phys. Chem. **29**, 403-452 (2010).


FTLM: J. Schnack and O. Wendland, Eur. Phys. J. B **78**, 535-541 (2010).

Icosidodecahedron $s = 1/2$



Exp. data: A. M. Todea, A. Merca, H. Bögge, T. Glaser, L. Engelhardt, R. Prozorov, M. Luban, A. Müller, Chem. Commun., 3351 (2009).

Summary



- One can indeed exploit $SU(2)$ and point group symmetries together. Good for molecules, since they are of finite size.
- **Problem: Recoupling coefficients.**
- Finite-temperature Lanczos is a good approximate method for Hilbert space dimensions smaller than 10^{10} .
- **I believe that this is the future.**

Many thanks to my collaborators worldwide

- T. Englisch, T. Glaser, M. Höck, N.B. Ivanov, S. Leiding, A. Müller, S. Ratnabala, R. Schnalle, Chr. Schröder, J. Ummethum, O. Wendland (Bielefeld)
- K. Bärwinkel, H.-J. Schmidt, M. Neumann (Osnabrück)
- M. Luban (Ames Lab, USA); P. Kögerler (Aachen, Jülich, Ames); R.E.P. Winpenny, E.J.L. McInnes (Man U, UK); L. Cronin, M. Murrie (Glasgow, UK); E. Brechin (Edinburgh, UK); H. Nojiri (Sendai, Japan); A. Postnikov (Metz, France)
- J. Richter, J. Schulenburg (Magdeburg); A. Honecker (Göttingen); U. Kortz (Bremen); A. Tennant, B. Lake (HMI Berlin); B. Büchner, V. Kataev, H.-H. Klauß (Dresden); P. Chaudhuri (Mühlheim); J. Wosnitza (Dresden-Rossendorf); J. van Slageren (Stuttgart); R. Klingeler (Heidelberg); O. Waldmann (Freiburg)

Thank you very much for your attention.

Molecular Magnetism Web

www.molmag.de

Highlights. Tutorials. Who is who. Conferences.